

WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: September 25, 2011 – October 8, 2011 (weeks 39 & 40)

Summary

In New Brunswick, low influenza activity for weeks 39 & 40 and within expected levels

New Brunswick:

- There have been no positive influenza detections during weeks 39 & 40, consistent with baseline levels.
- The ILI consultation rate in week 40 was higher than week 39 and slightly above the expected range for this time of year.
- No ILI or influenza outbreaks were reported in weeks 39 & 40, consistent with baseline levels.

Canada:

- Influenza activity continues at low inter-seasonal levels with few detections of influenza in weeks 39 & 40. Three regions (one each) in Quebec, British Columbia and Ontario reported sporadic influenza activity in weeks 39 & 40, similar to previous 2 weeks. An increase in detections of rhinovirus and parainfluenza was observed during the 2-week period.
- Six influenza detections were reported in weeks 39 & 40, similar to weeks 37 & 38.
- The ILI consultation rate was above the expected range for week 39. No outbreaks were reported in weeks 39 & 40.


Note: The weekly New Brunswick influenza reports will resume on October 24, 2011 (week 41).

1) Influenza Laboratory Data¹

- Overall influenza activity remains low in weeks 39 & 40.
- No influenza detections were reported during that period.
- Since the beginning of the season, no positive influenza detections have been reported.

Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of 7 sites in Emergency Rooms, 5 sites in Family Practice, 3 sites in First Nations communities, 1 site in a Nursing Home, 3 sites in Universities and 8 sites in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

<u>Graph 1</u>: Number and percent of positive influenza specimens in New Brunswick, by week, up to October 8, 2011 (data source: G. Dumont lab results)



<u>Table 1</u>: Positive influenza test results by Health Region in New Brunswick up to October 8, 2011 (data source: G. Dumont lab results)


Region	Reporting period: September/25/2011–October/08/2011						Cumulative: (2011/2012 season) Aug./28/2011 –Oct./08/2011				Cumulative: (2010/2011 season) Aug./29/2010 – Aug./27/2011			
	Activity level ²	А				В	A B			В	А		В	
		A(H1)	A(H3)	pH1N1	unsubt yped		A(H1)	A(H3)	pH1N1	unsubt yped		Non- pH1N1	pH1N1	
Region 1	No activity	0	0	0	0	0	0	0	0	0	0	433	56	29
Region 2	No activity	0	0	0	0	0	0	0	0	0	0	57	2	13
Region 3	No activity	0	0	0	0	0	0	0	0	0	0	112	16	19
Region 4	No activity	0	0	0	0	0	0	0	0	0	0	79	58	56
Region 5	No activity	0	0	0	0	0	0	0	0	0	0	26	3	1
Region 6	No activity	0	0	0	0	0	0	0	0	0	0	46	27	5
Region 7	No activity	0	0	0	0	0	0	0	0	0	0	32	3	3
Total NB		0	0	0	0	0	0	0	0	0	0	785	165	126

 $^{^2 \} Influenza\ activity\ level\ definition\ is\ available\ on\ the\ PHAC\ FluWatch\ website: \\ \underline{http://www.phac-aspc.gc.ca/fluwatch/11-12/def11-12-eng.php}$

2) ILI Consultation Rates³

- During week 40, the ILI consultation rate was 24.2 consultations per 1,000 patient visits, a higher rate than week 39 (12.5 consultations per 1,000 patient visits) and slightly above the expected levels for this time of year.
- During week 40, the sentinel response rate was 67% for the FluWatch sentinel physicians and 58% for the NB SPIN practitioners. (8 FluWatch and 11 NB SPIN)

<u>Graph 2</u>: ILI Consultation Rates in New Brunswick, by report week, season 2011/12 compared to previous seasons*



^{*} The mean rate was based on data from the 1996/97 to 2010/2011 seasons and excludes the Pandemic season (2009-2010).

³ A total of 31 practitioner sites (12 FluWatch sentinel physicians and 19 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

3) ILI and Laboratory-Confirmed Outbreak Data


<u>Table 2:</u> ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.

	Septembe	Reporting period: er/25/2011 –October/	Cumulative # of	Cumulative # of outbreaks		
	Lab-confirmed outbreaks in Nursing Homes*	Schools reporting ILI outbreaks**	Lab-confirmed outbreaks in Other Settings*	outbreaks season 2011-2012	season 2010-2011	
Region 1	0 out of 13	0 out of 74	0	0	17	
Region 2	0 out of 15	0 out of 81	0	0	21	
Region 3	0 out of 14	0 out of 95	0	0	12	
Region 4	0 out of 6	0 out of 22	0	0	12	
Region 5	0 out of 2	0 out of 18	0	0	17	
Region 6	0 out of 9	0 out of 35	0	0	10	
Region 7	0 out of 4	0 out of 27	0	0	22	
Total NB	0 out of 63	0 out of 352	0	0	111	

^{*}Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

^{**}Schools reporting greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.

<u>Graph 3</u>: Number of Influenza Outbreaks in Nursing Homes¹ and ILI Outbreaks in Schools² reported to Public Health in New Brunswick, by report week, season 2011/12.



¹ The National FluWatch definition of an outbreak in a nursing home is stated as two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

<u>National Flu Watch Program</u> - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: www.phac-aspc.gc.ca/fluwatch/index.html

Other Links:

World-

http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html Europe: http://www.euroflu.org/cgi-files/bulletin_v2.cgi and

http://www.ecdc.europa.eu/en/healthtopics/seasonal influenza/epidemiological data/Pages/Weekly Influenza Surveillance Overview.aspx

PAHO: http://new.paho.org/hq/index.php?option=com content&task=blogcategory&id=805&Itemid=569]

Australia: http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm]

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza weekly update.php

Argentina: http://www.msal.gov.ar/archivos/INFORME%20INFLUENZA%20PANDÉMICA%20 H1N1 %2005-08-

2009.pdf

South Africa: http://www.nicd.ac.za/
US: www.cdc.gov/flu/weekly/

² The National FluWatch definition of an ILI outbreak in a school is stated as absenteeism greater than 10% (or absenteeism that is higher (e.g.>5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.