

WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: April 8, 2012 – April 14, 2012 (week 15)

Summary

In New Brunswick, continued increase (since week 11) in percent positive lab results

New Brunswick:

- There have been 56 positive influenza detections during week 15, 1 influenza A (H3), 3 influenza A (H1N1) pdm09 and 52 influenza B.
- The ILI consultation rate was low and is within the expected range for this time of year.
- 2 new ILI school outbreaks were reported, both in Region 5.

Canada:

- The peak of activity for the 2011-2012 influenza season in Canada has passed as most indicators of influenza activity continue to decline. However, activity remains elevated in some regions of the country (i.e. Atlantic Region, Ontario, Alberta & British Columbia).
- 841 laboratory detections of influenza were reported, proportion of positive tests was 18.9%. To date this season, influenza B remains the predominant virus type circulating in provinces east of Saskatchewan.
- The ILI consultation rate was within the expected levels for this time of year. 44 influenza or ILI outbreaks were reported: 25 in long-term care facilities, 4 in hospitals and 15 in other settings.

International:


United States: On April 12, the CDC reported one new case of human infection with a novel influenza A virus in a child in Utah. The child was infected with an influenza A (H3N2) variant virus similar to those identified in the 12 human infections that occurred between July and November 2011. The child has recovered and contact with swine in the week preceding onset of the child's illness was reported. No such cases have been reported in Canada.

1) Influenza Laboratory Data¹

- Continued increase in percent positive lab results (since week 11).
- 56 influenza detections were reported during that period.
- Since the beginning of the season, 220 positive influenza detections have been reported; 27 were influenza A (H1N1)pdm09 (12%), 16 were influenza A (H3)(7%) and 177 were influenza B viruses(81%).

¹ Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of 7 sites in Emergency Rooms, 5 sites in Family Practice, 3 sites in First Nations communities, 1 site in a Nursing Home, 3 sites in Universities and 8 sites in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

<u>Graph 1</u>: Number and percent of positive influenza specimens in New Brunswick, by week, up to April 14, 2012 (data source: G. Dumont lab results)



<u>Table 1</u>: Positive influenza test results by Health Region in New Brunswick up to April 14, 2012 (data source: G. Dumont lab results)


Region	Reporting period: April/08/2012-April /14/2012						Cumulative: (2011/2012 season) Aug./28/2011 –Apr./14/2012				Cumulative: (2010/2011 season) Aug./29/2010 – Aug./27/2011			
	Activity	А				В	A B				В	Α		В
	level ²	A(H1)	A(H3)	(H1N1) pdm09	unsubt yped		A(H1)	A(H3)	(H1N1) pdm09	unsubt yped		Non- pH1N1	pH1N1	
Region 1	Sporadic	0	1	0	0	43	0	13	14	0	157	433	56	29
Region 2	Sporadic	0	0	0	0	4	0	1	2	0	5	57	2	13
Region 3	No activity	0	0	0	0	0	0	1	0	0	1	112	16	19
Region 4	Sporadic	0	0	2	0	0	0	1	5	0	1	79	58	56
Region 5	No activity	0	0	0	0	0	0	0	1	0	1	26	3	1
Region 6	Sporadic	0	0	1	0	3	0	0	4	0	8	46	27	5
Region 7	Sporadic	0	0	0	0	2	0	0	1	0	4	32	3	3
Total NB		0	1	3	0	52	0	16	27	0	177	785	165	126

² Influenza activity level definition is available on the PHAC FluWatch website: http://www.phac-aspc.gc.ca/fluwatch/11-12/def11-12-eng.php

2) ILI Consultation Rates³

- During week 15, the ILI consultation rate was 21.5 consultations per 1,000 patient visits, and is within the expected levels for this time of year.
- During week 15, the sentinel response rate was 36% for the FluWatch sentinel physicians and 33% for the NB SPIN practitioners. (5 FluWatch and 6 NB SPIN)

<u>Graph 2</u>: ILI Consultation Rates in New Brunswick, by report week, season 2011/12 compared to previous seasons*



^{*} The mean rate was based on data from the 1996/97 to 2010/2011 seasons and excludes the Pandemic season (2009-2010).

³ A total of 32 practitioner sites (14 FluWatch sentinel physicians and 18 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

3) ILI and Laboratory-Confirmed Outbreak Data


<u>Table 2:</u> ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.

	Арг	Reporting period: il/08/2012 –April/14/	Cumulative # of	Cumulative # of outbreaks		
	Lab-confirmed outbreaks in Nursing Homes*	Schools reporting ILI outbreaks**	Lab-confirmed outbreaks in Other Settings*	outbreaks season 2011-2012	season 2010-2011	
Region 1	0 out of 13	0 out of 74	0	2	17	
Region 2	0 out of 15	0 out of 81	0	1	21	
Region 3	0 out of 14	0 out of 95	0	8	12	
Region 4	0 out of 6	0 out of 22	0	2	12	
Region 5	0 out of 2	2 out of 18	0	6	17	
Region 6	0 out of 9	0 out of 35	0	2	10	
Region 7	0 out of 4	0 out of 27	0	1	22	
Total NB	0 out of 63	2 out of 352	0	22	111	

^{*}Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

^{**}Schools reporting greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.

<u>Graph 3</u>: Number of Influenza Outbreaks in Nursing Homes¹ and ILI Outbreaks in Schools² reported to Public Health in New Brunswick, by report week, season 2011/12.



¹ The National FluWatch definition of an outbreak in a nursing home is stated as two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

<u>National Flu Watch Program</u> - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: <u>www.phac-aspc.gc.ca/fluwatch/index.html</u>

Other Links:

World-

http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html Europe: http://www.euroflu.org/cgi-files/bulletin_v2.cgi and

http://www.ecdc.europa.eu/en/healthtopics/seasonal_influenza/epidemiological_data/Pages/Weekly_Influenza_Surveillance Overview.aspx

PAHO: http://new.paho.org/hq/index.php?option=com_content&task=blogcategory&id=805&Itemid=569]

Australia: http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm]

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza weekly update.php

Argentina: http://www.msal.gov.ar/archivos/INFORME%20INFLUENZA%20PANDÉMICA%20_H1N1_%2005-08-2009.pdf

South Africa: http://www.nicd.ac.za/
US: www.cdc.gov/flu/weekly/

² The National FluWatch definition of an ILI outbreak in a school is stated as absenteeism greater than 10% (or absenteeism that is higher (e.g.>5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.